

# The complete sequence of the mitochondrial genome of the Chinook salmon, *Oncorhynchus tshawytscha*.

VIVIAN WILHELM<sup>1,2</sup>, JAIME VILLEGAS<sup>1,2</sup>, ÁLVARO MIQUEL<sup>3</sup>, ESTEBAN ENGEL<sup>1,2</sup>, SEBASTIÁN BERNALES<sup>1,2</sup>, PABLO D.T. VALENZUELA<sup>1,2</sup> AND LUIS O. BURZIO<sup>1,2,3</sup>

<sup>1</sup>Fundación Ciencia para la Vida, <sup>2</sup>Instituto Milenio de Biología Fundamental y Aplicada, MIFAB, and <sup>3</sup>Bios Chile I.G.S.A. Avenida Marathon 1943, Santiago, Chile

## ABSTRACT

The complete sequence of the mitochondrial genome of Chinook salmon, *Oncorhynchus tshawytscha*, has been determined. The circular genome consisting of 16,644 base pairs encodes thirteen proteins, the 12S and 16S ribosomal RNAs, and 22 transfer RNAs. These genes are ordered in the same way as most other vertebrates. The nucleotide and amino acid sequences of the ribosomal RNAs and the thirteen protein-coding genes were compared with those of other salmonids such as *Oncorhynchus mykiss*, *Salmo salar*, *Salvelinus fontinalis*, *Salvelinus alpinus* and *Coregonus lavaretus*. The sequence features of the control region (D-loop), the origin of L-strand replication and a putative peptide codified by the 16S mitochondrial RNA are described and discussed.

**Key terms:** Salmonids, Mitochondria, Genome, Peptides, 16S rRNA

## INTRODUCTION

Since the publication of the complete sequence of the human mitochondrial DNA (mtDNA) by Anderson *et al.*, (1981), the organization of the mitochondrial genome of a variety of organisms including multicellular animals, plants, fungi and protozoa has been well characterized (Wolstenholme, 1992). At present, the mtDNA of 67 fish have been sequenced (NCBI: [www.ncbi.nlm.nih.gov/PMGifs/Genomes/7898.html](http://www.ncbi.nlm.nih.gov/PMGifs/Genomes/7898.html)). Due to the relative lack of recombination of the mtDNA and its maternal mode of inheritance, the mitochondrial genome represents a useful marker to use in population and phylogenetic studies. In this regard, the salmonid fish are commercially important, and the farming of these fish represents a major economic activity in Norway, Chile, Canada, Scotland, and elsewhere. Consequently, there is great interest in knowing the genetic features of the

mitochondrial DNA for stock identification, management, conservation, and population studies (Phillips and Oakley, 1997). Within this genus the complete sequence of the mtDNA of the rainbow trout *O. mykiss* (Zardoya *et al.*, 1995) was followed by the sequence of the mtDNA of Atlantic salmon *S. salar* (Hurst *et al.*, 1999), and other salmonids such as *S. alpinus* (Doiron *et al.*, 1999), *C. lavaretus* (Miya and Nishida, 2000) and *S. fontinalis* (Doiron *et al.*, 2002). Similar to other vertebrates, the organization of the mtDNA of salmonid fish species contains 13 protein coding genes, 22 transfer tRNA genes, and 2 ribosomal RNAs genes corresponding to the 12S and 16S transcripts.

Recently we have been involved in sequencing the complete genome of *Piscirickettsia salmonis* (Valenzuela *et al.*, 2001), the causative agent of SRS, a disease with negative consequences for the salmon farming industry (Fryer and Muel, 1997). *P. salmonis* is a Gram-negative obligate

intracellular bacterium found in cytoplasmic vacuoles of infected cells. To grow the pathogen, fish cells such as CHSE-214 obtained from Chinook salmon (*O. tshawytscha*) (Lannan *et al.*, 1984) are infected with the bacteria, and after 10 to 12 days, *P. salmonis* is harvested (Jamett *et al.*, 2001). To achieve the purification of the bacterial DNA and to eliminate contamination from CHSE-214 cell DNA, we developed a procedure based on the treatment with DNase I. To avoid the alteration of the bacterial membranes and the consequent destruction of the *P. salmonis* DNA, we developed a buffer solution to preserve the integrity of the bacterial cells. This condition involves the suspension of the purified microorganisms in a solution that also stabilized the mitochondria of the CHSE-214 cells. Therefore our final preparation of DNA of *P. salmonis* was contaminated with the mitochondrial DNA of Chinook salmon. We accepted this risk as the low complexity of the mitochondria genome would not alter the assembly of the different fragments of the shotgun library into contigs. Here we present the complete sequence of the mtDNA of Chinook salmon based on 8 contigs that were completed by PCR amplification to close the gaps. The sequence features of the control region (D-loop), the origin of replication of the L-strand replication, and a putative peptide codified by the 16S mitochondrial RNA, are discussed.

#### MATERIALS AND METHODS

##### *Growth of P. salmonis in Chinook salmon cells.*

*P. salmonis* LF-89 (ATCC VR 1361) were continuously propagated in Chinook salmon embryo cell line CHSE-214 (CRL 1681 ATCC) (Lannan *et al.* 1984) using the antibiotic-free Minimal Essential Medium (MEM) plus 5% bovine serum as described elsewhere (Jamett *et al.* 2001). To follow the growth of the pathogen, the amount of bacteria in the culture medium was evaluated by using an ELISA test for *P. salmonis* (Aguayo *et al.*, 2002).

##### *Purification of P. salmonis and Chinook salmon mitochondria.*

When the cytopathic effect on the CHSE-214 was more than 90%, the lysate was collected from the culture flasks and centrifuged at 200xg. The pellet containing *P. salmonis* and cellular debris was resuspended in a modified PBS solution containing 200 mM NaCl (Jamett *et al.*, 2001) and centrifuged at 200 x g. This washing procedure was repeated once, and the final pellet, referred to as the partially-purified fraction of *P. salmonis* (Jamett *et al.* 2001), was resuspended in 5 ml of the PBS-200 mM NaCl solution. To determine the number of bacteria per ml, serial dilutions in the PBS-200 mM NaCl solution were counted under phase microscopy in a Neubauer chamber. Using this procedure, one mg of protein corresponds to approximately  $3.8 \times 10^9$  *P. salmonis* (Jamett *et al.* 2001).

##### *DNA isolation*

The partially-purified *P. salmonis* fraction was resuspended in 5 ml of a solution containing 50 mM Tris-HCl, pH7.5, 200 mM NaCl, 2mM KCl, 5 mM MgCl<sub>2</sub>, 0.1 mM CaCl<sub>2</sub>, 2 mM 2-mercaptoethanol, 2 µg/ml of aprotinin and 2 µg/ml of leupeptin. To this mixture, 350 µg of Dnase I and 350 µg of Rnase A were added and then incubated at 25°C for 60 min. To stop the reaction, 20 volumes of PBS containing 200 mM NaCl and 1mM EDTA were added, and the mixture was centrifuged at 7,000 x g for 20 min. The bacterial pellet was washed once more with the above PBS solution and the final pellet was resuspended in 5 ml of the lysis buffer (50 mM Tris-HCl, pH 9.0, 400 mM NaCl, 20 mM EDTA, 0.5 M sucrose and 0.2% SDS). To the suspension, 1 mg of proteinase K was added, and the mixture was incubated at 58°C for 60 min. The DNA was then extracted with the phenol procedure as described (Sambrook *et al.*, 1989). To check for the absence of nuclear salmon DNA, the preparation was used as a template to amplify the 18S ribosomal gene (forward: 5' ACGGCCGGTACAGTGAAACT; reverse:

5' CCAATTACAGGGCCTCGAAA) (Villegas *et al.*, 2000). Those preparations of the bacteria DNA without contamination with salmon DNA were used for sequencing.

#### DNA sequencing

The mixture of bacterial and mitochondrial DNA purified as described was sequenced by the whole-genome shotgun approach as described before for the complete sequence of several bacteria (Stover *et al.*, 2000). The DNA was fragmented by nebulization to an average size of approximately 1,000 bp. The random fragments were cloned into the pIK96 plasmid using the *Bst*XI adaptors, provided by Incyte Genomics Inc. A total of 18,887 sequence reads were obtained during the random sequencing phase using ABI 3700 sequencers at Incyte Genomics Inc. (Palo Alto, CA) and ABI 310 DNA sequencers in our laboratory.

#### Bioinformatics

The sequences of the fragments were assembled and edited into contigs using the SEQUENCHER 4.1 and the Vector NTI 6.1 programs. The gaps between contigs corresponding to the mitochondrial DNA were closed and sequenced by PCR (Villegas *et al.*, 2002) and with primers designed according to the sequence adjacent to the 3' and 5' ends of the corresponding contigs. The mitochondrial genes were identified using the sequence-alignment algorithms BLASTX and BLASTP (Altschul *et al.*, 1997) and the complete mitochondrial genome of the salmonids *O. mykiss* (NC\_001717), *S. salar* (U12143), *S. alpinus* (AF154851), *S. fontinalis* (NC\_000860) and *C. lavaretus* (AB034824).

## RESULTS AND DISCUSSION

#### Genome Organization


As described in Methods, a fraction of the DNA preparation of *P. salmonis*

corresponded to the mtDNA of CHSE-214 cells used to grow the bacteria (Jamett *et al.*, 2001). Within the 2,143 contigs assembled from 18,887 reads of the bacterium genome, we identify 8 contigs that correspond to the mtDNA of *O. tshawytscha*. The 9 gaps were closed by PCR, followed by a cloning of the corresponding amplicon and sequencing. The complete sequence of *O. tshawytscha* contains 16,644 bp, and it has been deposited in the NCBI/GeneBank libraries under the accession number NC\_002980.

To date, the complete sequence of 67 mitochondrial genomes of fish have been deposited in the GeneBank ([www.ncbi.nlm.nih.gov/PMGifs/Genomes/7898.html](http://www.ncbi.nlm.nih.gov/PMGifs/Genomes/7898.html)), including the mtDNA of six salmonid species. The size of the mtDNA of *O. tshawytscha* is similar to those of other salmonids such as *O. mykiss* (16,642 bp), *S. salar* (16,665 bp), *S. fontinalis* (16,624 bp), *S. alpine* (16,659 bp) and *C. lavaretus* (16,737 bp). The genome comprises 2 ribosomal genes, 13 protein-encoding genes, 22 tRNAs, and the control region. The position and orientation of the mitochondrial genes and the control region are similar to those described in most other fish and higher vertebrates (Fig. 1). The overlapping arrangement between the set of genes *ATP8/ATP6* and *ND4L/ND4* were 10 and 7 bp, respectively. This overlapping is identical to that found in *O. mykiss*, *S. salar*, *S. alpinus*, *S. fontinalis* and *C. lavaretus*. Although the genes of *ND5* and *ND6* are encoded in the H and L-strand, respectively, the overlapping is also similar to that found in the other salmonids.

#### Protein-Encoding Genes

The Chinook salmon mtDNA contains 13 open reading frames, 12 codified by the H-strand and one (*ND6*) codified by the L-strand (Fig. 1). In general, there is a high homology of sequences at the nucleotide and amino acid level when they are compared with other salmonid species (Table I). All the initiation codons are ATG except that of *COI* gene, which is GTG. This initiation codon also has been described in other fish

**Figure 1**

The organization of the mitochondrial genome of *O. tshawytscha*. The outer and the inner circles represent the H- and L-strands, respectively. The tRNA genes are indicated by the single letter amino acid code.

TABLE I

Percentage of nucleotide (Nt) and amino acid (AA) sequence identities of *O. tshawytscha* mitochondrial genes against their homologues of other salmonids.

Genes	<i>O. mykiss</i>		<i>S. salar</i>		<i>S. fontinalis</i>		<i>S. alpinus</i>		<i>C. lavaretus</i>	
	Nt	AA	Nt	AA	Nt	AA	Nt	AA	Nt	AA
12S rRNA	97.3	-	96.0	-	96.3	-	96.5	-	94.9	-
16S rRNA	96.4	-	95.6	-	95.7	-	95.8	-	94.7	-
NAD 1	91.6	100	86.1	96.9	85.4	97.5	87.0	98.1	82.4	96.9
NAD 2	90.2	95.4	86.0	94.0	85.6	92.8	85.6	92.8	78.6	87.1
COI	93.8	99.8	87.8	99.0	89.5	99.4	89.6	99.4	85.2	99.0
COII	94.1	98.3	92.5	97.8	90.8	97.8	91.0	97.8	88.0	97.4
ATP8	96.4	100	98.2	100	95.2	98.1	95.8	96.4	95.8	100
ATP6	89.0	97.8	87.4	97.8	85.6	98.2	85.7	98.2	83.6	97.8
COIII	86.2	95.8	89.9	97.3	88.5	96.5	90.3	96.9	86.2	95.8
NAD 3	92.5	99.1	84.5	94.0	88.5	95.7	86.5	94.0	79.3	93.2
NAD 4L	94.6	100	91.2	99.0	89.8	100	89.6	100	88.5	100
NAD 4	91.3	97.8	86.9	97.2	85.9	96.7	87.5	96.5	83.0	97.8
NAD 5	92.4	96.9	87.6	94.3	87.0	93.4	87.6	94.0	82.5	91.3
NAD 6	92.3	98.8	86.0	94.2	86.9	95.3	86.0	96.5	81.8	95.3
Cyt b	92.1	99.2	87.8	98.7	88.1	98.4	87.8	98.4	84.5	98.4

species including the other salmonids described before. On the other hand, the termination codon TAA is present in the ORF of *COI*, *ATP8*, *ATP6*, *COIII*, *ND4L* and *ND5*, whereas codon TAG is present in *ND1*, *ND2* and *ND6*. In the later gene, this termination codon is present in all the mtDNA of salmonids species so far sequenced. For the other genes, the TAG termination codon varies among the species. The four remaining genes (*COXII*, *ND3*, *ND4*, *Cyt b*) do not end with a complete stop codon but with single T. In humans (Ojala *et al.*, 1981) and in *Xenopus* (Roe *et al.*, 1985) it was demonstrated that the transcript of these gene contains a complete termination codon TAA created by post-transcriptional polyadenylation. It is reasonable to argue that a similar situation takes place after transcription of these mitochondrial genes in *O. tshawytscha* as well as in other salmonids that also contain incomplete TAA termination codon in their mitochondrial genome, as suggested previously (Zardoya *et al.*, 1995; Hurst *et al.*, 1999).

#### Ribosomal and transfer RNA

The 12S and 16S rRNA genes in *O. tshawytscha* mtDNA are 947 and 1,683 bp long, respectively. Similar to other vertebrates, the rRNA genes are located between tRNA<sup>Phe</sup> and tRNA<sup>Leu</sup> (UUR) and

are separated by tRNA<sup>Val</sup> (Fig. 1). Alignment of the ribosomal genes with those of other salmonids revealed high identity values, indicating that they are highly conserved among these species (Table I). On the other hand, alignment with the corresponding genes of other fish and mammals revealed a typical pattern of conserved segments interrupted by variable sequences (data not shown).

An interesting feature found in the 16S rRNA is the presence of short ORFs as shown in Figure 2, using the vertebrate mitochondrial code. If the sequences of the salmonids 16S rRNAs are used to run any program for nucleic acid translation, several peptides of different length are identified and only a few of them began with Met. Therefore, the only ORFs selected were those that corresponded to a minimum of 24 amino acids, present in approximately the same position in the 16S rRNA and which show high amino acid homology among the six salmonids species considered in this study. In *O. tshawytscha* an ORF for a peptide of 46 amino acids and codified by the H-strand between positions 976 and 1,113 was found (Fig. 2). The amino acid sequence of this peptide shows a 97.8% of identity with similar peptides found in the other salmonid species and with similar position in the 16S rRNA (Fig. 2). An ORF corresponding to a second peptide around position 1223 and 1377 of the H-strand of the 16S rRNA was also found. However, this

<b><i>O. tshawytscha</i></b>	<b>MGLTAAVFWPCEGSAITCLLNEDLYEWHHEGLAVSSFKSMKLICPC</b>	<b>(976 - 1113)</b>
<b><i>S. salar</i></b>	<b>MGLTAAVFWPCEGSAITCLLNEDLYEWHHEGLAVSSSKSMKLICPC</b>	<b>(973 - 1110)</b>
<b><i>O. mykiss</i></b>	<b>MGLTAAVFWPCEGSAITCLLNEDLYEWHHEGLAVSSSKSMKLICPC</b>	<b>(970 - 1107)</b>
<b><i>S. fontinalis</i></b>	<b>MGLTAAVFWPCEGSAITCLLNEDLYEWHHEGLAVSSPKSMKLICPC</b>	<b>(973 - 1110)</b>
<b><i>S. alpinus</i></b>	<b>MGLTAAVFWPCEGSAITCLLNEDLYEWHHEGLAVSSPKSMKLICPC</b>	<b>(973 - 1110)</b>
<b><i>C. lavaretus</i></b>	<b>MGLTAAVFWPCEGSAITCLLNEDLYEWHHEGLAVSSSKSVKLICPC</b>	<b>(971 - 1108)</b>

**Figure 2**

Sequences of the peptide-coding ORF in the 16S ribosomal RNA of the *O. tshawytscha* and five other salmonids. The numbers in parenthesis at the right of each sequence correspond to the position in the 16S rRNA.

peptide shows less identity between the salmonids and variation in the length between 49 (*S. salar*) and 60 (*O. mykiss*) amino acids (data not shown). In addition to the restriction imposed on selecting the peptide, it is reasonable to ask whether they have a physiological meaning. This search was inspired by recent work that described a peptide of 24 amino acids, referred to as humanin, a factor that protects the neuronal cell from the apoptosis that occurs in familial Alzheimer's disease (Niikura *et al.*, 2002). The human 16S mitochondrial gene codifies this peptide and the ORF is in positions 963 to 1037, similar to the positions where we found the ORF in the salmonid ribosomal RNA (Fig. 2). Therefore, it might be possible that these putative peptides play a functional role in salmonids. This finding might be another surprise of the pleiotrophic features of the 16S mitochondrial RNA in addition to other intriguing properties such as nuclear localization (Villegas *et al.*, 2000; 2002), cytoplasmic localization, the role in the development of the germ cell line (Kobayashi *et al.*, 1998), and the chaperone-like activity (Sulijoadikusumo *et al.*, 2001).

#### tRNA sequences

Similar to other vertebrates, the mtDNA of *O. tshawytscha* contains 22 tRNA genes interspersed between ribosomal RNAs and protein encoding regions. Fourteen tRNAs are codified by the H-strand and 8 are codified by the L-strand, similar to the organization in other salmonids. As reported before, the tRNA<sup>Met</sup> is quite well conserved, showing high identity with the sequence found in *S. salar*, *O. mykiss*, *S. alpinus*, *S. fontinalis*, *C. lavaretus* (data not shown). The DHU loop of tRNA<sup>Leu</sup> (UUR) is highly conserved among vertebrates and the sequence of this loop shows high identity with those of *O. mykiss*, *S. salar* and other salmonids (data not shown). The sequence of this loop has been proposed as necessary for the correct stop of transcription of the ribosomal RNAs plus the included tRNAs (Shadel and Clayton, 1997). This is an important control region

of transcription of the mtDNA to allow the synthesis of as many copies of the rRNAs needed for translation (Shadel and Clayton, 1997; Taanman, 1999). All the tRNAs of *O. tshawytscha* can be folded into the cloverleaf secondary structures (data not shown), and this folding requires the formation of G-U pairing as reported before (Wolstenholme, 1992).

In contrast with the other tRNAs, the tRNA<sup>Ser</sup> (AGY) contains a complete DHU arm, which is missing in the rest of the homologous tRNAs and as proposed previously, this is an exception conserved in fish (Wolstenholme, 1992).

#### The Control Region

Between the tRNA<sup>Pro</sup> and tRNA<sup>Phe</sup> genes of vertebrates mtDNAs there is a non-coding region that varies in length (Wolstenholme, 1992). Because in mammals and amphibians this sequence has been shown to include the signals required for the initiation of the H-strand replication, and it has been designated as the control region for both H-strand and L-strand transcription (Shadel and Clayton, 1997). In *O. tshawytscha* the control region is 986 bp, similar to the size of *S. salar* (1,006 bp), *O. mykiss* (1003 bp), *S. fontinalis* (964 bp), *S. alpinus* (998 bp) and *C. lavaretus* (1,076 bp).

Adjacent to the tRNA<sup>Pro</sup> gene it is possible to find in the D-loop region the putative TAS sequences associated with premature termination of the replication cycle (Taanman, 1999), the polypirimidine track that varies between 24 to 25 T and C and that could be a site of interaction with the mitochondrial single-strand-DNA-binding protein (Mignotte *et al.*, 1985), and the imperfect direct repeat found in the right domain close to the tRNA<sup>Phe</sup>, region that harbor the transcription promoters in human and mouse (Clayton 1992; Taanman 1999). The CBS motifs CSB-2 and CSB-3 near the origin of replication are highly conserved in the six salmonid species studied as shown in Figure 3. At approximately the position expected for the CSB-1 motive, relative to CSB-2, the sequence 5' ACATA is found in all the fish,

CSB-2		CSB-3	
<i>O. tshawytscha</i>	TAAACCCCCCTACCCCCCT	<i>O. tshawytscha</i>	TGTTAAACCCCTAAACCAG
<i>O. mykiss</i>	TAAACCCCCCTACCCCCCT	<i>O. mykiss</i>	TGTTAAACCCCTAAACCAG
<i>S. salar</i>	CAAACCCCCCTACCCCCCT	<i>S. salar</i>	TGTCAAACCCCTAAACCAG
<i>S. alpinus</i>	TAAACCCCCCTACCCCCCT	<i>S. alpinus</i>	TGTTAAACCCCTAAACCAG
<i>S. fontinalis</i>	TAAACCCCCCTACCCCCCT	<i>S. fontinalis</i>	TGTTAAACCCCTAAACCAG
<i>C. lavaretus</i>	TAAACCCCCCTACCCCCCT	<i>C. lavaretus</i> 7	TGTCAAACCCCAAACCAG

**Figure 3**

Comparative alignment of CSB motifs identified in the control region of the mtDNA of *O. tshawytscha* and five other salmonids species.

which is an imperfect motif if compared with the CSB-1 of mouse and human (Walberg and Clayton 1981; Shadel and Clayton 1997) and already discussed by Digby *et al.*, (1992).

It was reported that in the left domain of the control region of *O. mykiss* there is an ORF corresponding to 102 codons (Digby *et al.*, 1992). However, we were unable to find similar coding region in the control region of the other salmonids.

#### *The L-strand replication*


It has been described previously that in the mtDNA of several species there is a putative origin of L-strand replication in a cluster of five tRNAs genes (WANCY region) and between the tRNA<sup>Asn</sup> and tRNA<sup>Cys</sup>. This sequence has the potential to fold in a stem-loop secondary structure with several characteristics. In human mtDNA the 5'-GCCGG-3' motif is involved in the transition from RNA to DNA synthesis (Hixson *et al.*, 1986), and the same sequence was found in this region of *O. tshawytscha* as well as in other salmonids species (Fig. 4). The stem sequence is conserved among vertebrates, whereas the loop sequence is more variable (Fig. 4). It is interesting that

the loop of *O. tshawytscha* is the shortest one (11 nucleotides) compared with the longest loop of *O. mykiss* of 17 nucleotides. In all the mammalian mtDNA already sequenced the loop is a T-rich sequence that was proposed to be necessary for the initiation of the L-strand replication by a RNA primase (Taanman, 1999). This is in contrast with the loop of *O. tshawytscha* which is C-rich similar to other salmonids (Fig. 4). Therefore, it seems that the initiation of L-strand replication is not restricted to a stretch of T as previously suggested (Taanman, 1999) but rather to a stretch of polypirimidine.

The sequence in the control region close to the gene of tRNA<sup>Pro</sup> gene, shows great variation among these important species for the salmon industry, characteristic that offers an important mtDNA markers to be used to study the *O. tshawytscha* population and for the certification of eggs and smolts.

#### ACKNOWLEDGEMENTS

We are grateful to Dr. Mario Roseblatt and Arturo Yudelevich for helpful discussion during the course of this research and critical reading of this manuscript.



**Figure 4**

Theoretical secondary structure for the replication origin of the L-strand of *O. tshawytscha*. The structure is compared with that of *S. salar*, *O. mykiss*, *S. alpinus*, *S. fontinalis* and *C. lavaretus*. The nucleotide sequences of the H-strand template are shown. The sequence associated with the transition between RNA synthesis and DNA synthesis is underlined.

#### REFERENCES

- AGUAYO J, MIQUEL A, ARANKI N, JAMETT A, VALENZUELA PDT, BURZIO LO (2002) Detection of *Piscirickettsia salmonis* in fish tissues by an enzyme-linked immunosorbent assay using specific monoclonal antibodies. *Dis Aquat Org* 49: 33-38
- ALTSCHUL SF, MADDEN TL, SCHAFFER AA, ZHANG J, ZHANG Z, MILLER W, LIPMAN DJ (1997) Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Res* 25: 3389-3402
- ANDERSON S, BANKIER AT, BARRELL BG, DE BRUIJN MH, COULSON AR, DROUIN J, EPERON IC, NIERLICH DP, ROE BA, SANGER F, SCHREIER PH, SMITH AJ, STADEN R, YOUNG IG (1981) Sequence and organization of the human mitochondrial genome. *Nature* 290: 457-465
- CLAYTON DA (1992) Transcription and replication of animal mitochondrial DNAs. *Int. Rev. Cytol.* 141: 217-232
- DIGBY TJ, GRAY MW, LAZIER CB (1992) Rainbow trout mitochondrial DNA: sequence and structural characteristics of the non-coding control region and flanking tRNA genes. *Gene* 118, 197-204
- DOIRON S, BLIER PU, BERNATCHEZ L (1999) NCBI Access N° AF154851
- DOIRON S, BLIER PU, BERNATCHEZ L (2002) NCBI Access N° NC\_000860
- FRYER JL, MAUEL MJ (1997) The Rickettsia: an emerging group of pathogens in fish. *Emerg. Infect. Dis.* 3: 137-144
- HIXSON JE, WONG TW, CLAYTON DA (1986) Both the conserved stem-loop and divergent 5'-flanking sequences are required for initiation at the human mitochondrial origin of light-strand DNA replication. *J Biol Chem* 261: 2384-2390



- HURST CD, BARTLETT SE, DAVIDSON WS, BRUCE IJ, (1999) The complete nucleotide sequence of the mitochondrial DNA of the Atlantic salmon, *Salmo salar*. *Gene* 239: 237-242
- JAMETT A, AGUAYO J, MIQUEL A, MULLER I, ARRIAGADA R, BECKER MI, VALENZUELA PDT, BURZIO LO (2001) Characterization of monoclonal antibodies against *Piscirickettsia salmonis*. *J Fish Dis* 24: 205-215
- KOBAYASHI S, AMIKURA R, MUKAI M (1998) Localization of mitochondrial large ribosomal RNA in germ plasm of *Xenopus* embryos. *Current Biol* 8: 1117-1120
- LANNAN CN, WINTON JR, FRYER JL (1984) Fish cell lines: establishment and characterization of nine cell lines from salmonids. *In Vitro* 20:671-676
- MIGNOTTE B, BARAT M, MOUNOLOU JC (1985) Characterization of a mitochondrial protein binding to single-stranded DNA. *Nucleic Acids Res* 13: 1703-1716
- MIYA M, NISHIDA M (2000) Use of Mitogenomic Information in Teleostean Molecular Phylogenetics: A Tree-Based Exploration under the Maximum-Parsimony Optimality Criterion. *Mol Phylogenet Evol* 17: 437-455
- NIKURA T, HASHIMOTO Y, TAJIMA H, NISHIMOTO I. (2002) Death and survival of neuronal cells exposed to Alzheimer's insults. *J Neurosci Res* 70: 380-391
- OJALA D, MONTOYA J, ATTARDI G (1981) tRNA punctuation model of RNA processing in human mitochondria. *Nature* 290: 470-474
- PHILLIPS RB, OAKLEY TH (1997) Phylogenetic relationships among the Salmonidae based on nuclear DNA and mitochondrial DNA sequences. In: KOCHER T, STEPIEN C (eds) *Molecular Systematics of Fishes*. San Diego: Academic Press. pp:145-162
- ROE BA, MA DP, WILSON RK, WONG JF (1985) The complete nucleotide sequence of the *Xenopus laevis* mitochondrial genome. *J Biol Chem* 260: 9759-9774
- SAMBROOK J, FRITSCH E, MANIATIS T (1989) *Molecular Cloning, A Laboratory Manual*. 2<sup>nd</sup> ed, New York: Cold Spring Harbor Laboratory Press
- SHADEL GS, CLAYTON DA (1997) Mitochondrial DNA maintenance in vertebrates. *Ann Rev Biochem* 66: 409-435
- STOVER CK, PHAM XQ, ERWIN AL, MIZOGUCHI SD, WARRENER P, HICKEY MJ, BRINKMAN FS, HUFNAGLE WO, KOWALIK DJ, LAGROU M, GARBER RL, GOLTRY L, TOLENTINO E, WESTBROCK-WADMAN S, YUAN Y, BRODY LL, COULTER SN, FOLGER KR, KAS A, LARBIG K, LIM R, SMITH K, SPENCER D, WONG GK, WU Z, PAULSEN IT, REIZER J, SAIER MH, HANCOCK RE, LORY S, OLSON MV (2000) Complete genome sequence of *Pseudomonas aeruginosa* PA01, an opportunistic pathogen. *Nature* 406: 959-964
- SULIJOADIKUSUMO I, HORIKOSHI N, USHEVA A (2001) Another function for the mitochondrial ribosomal RNA: protein folding. *Biochem* 40: 11559-11564
- TAANMAN JW (1999) The mitochondrial genome: structure, transcription, translation and replication. *Biochim Biophys Acta* 1410: 103-123
- VALENZUELA PDT, BURZIO LO, ROSEMBLATT M, YUDELEVICH A, BERNALES, S, ENGEL E, ERAZO E, HERNÁNDEZ C, HUARACAN B, ARAYA P, MARTÍNEZ R, MIQUEL A, MORALES C, SOZA C, VILLEGAS J, WILHELM V (2001) Sequence and applications of the *Piscirickettsia salmonis* genome. *Biol Res* 34: R17
- VILLEGAS J, ZARRAGA AM, MULLER I, MONTECINOS L, WERNER E, BRITO M, MENESES AM, BURZIO LO (2000) A novel chimeric mitochondrial RNA localized in the nucleus of mouse sperm. *DNA Cell Biol* 19: 579-588
- VILLEGAS J, ARAYA P, BUSTOS-OBREGON E, BURZIO LO (2002) Localization of the 16S mitochondrial rRNA in the nucleus of mammalian spermatogenic cells. *Mol Hum Reprod* 8: 977-983
- WALBERG MW, CLAYTON DA (1981) Sequence and properties of the human KB cell and mouse L cell D-loop regions of mitochondrial DNA. *Nucleic Acids Res* 9: 5411-5421
- WOLSTENHOLME DR (1992) Animal mitochondrial DNA: structure and evolution. *Int Rev Cytol* 141:173-216
- ZARDOYA R, GARRIDO-PERTIERRA A, BAUTISTA JM (1995) The complete nucleotide sequence of the mitochondrial DNA genome of the rainbow trout, *Oncorhynchus mykiss*. *J Mol Evol* 41: 942-951